



Ethnoecological knowledge associated to wild medicinal plants in district lower Dir Khyber Pakhtun Khwa Pakistan

Shakir Ullah¹, Zakir Uallah², Javeed Iqbal³, Fozia Abasi⁴, Siraj Khan⁵, Mohammad Sohail⁶, Mohsin Ihsan⁷

¹⁻⁴ Department of Botany, Govt Post Graduate Collage Timargara, Lower Dir, Pakistan

^{5,6} Abdul Wali Khan University, Department of Botany Garden Campus, Mardan, Pakistan

⁷ Department of Botany, Women University of Azad Jammu and Kashmir, Bagh Pakistan

Abstract

The present study is initiated to assess and document the existing ethnoecological knowledge of the indigenous people of District Lower Dir. For this purpose, 27 villages were chosen using purposive sampling methods based on the availability of main informants. Both purposively and randomly informants age ranged from 19 to 86 were chosen from among those born or have lived there. Quantitative and qualitative ethnobotanical methods were used to analyze the ethnological data. Significance test on the indigenous knowledge variation of the average number of reported medicinal plants were assessed. Result indicated that a total of 112 wild medicinal plants belonging to 53 families were recorded from the study sites. Moreover, the indigenous knowledge about the medicinality of the reported medicinal plants were found to be evenly known by all informants regardless of their gender, age, level of education, marital status, and experiences. On top of this, it was found that the District has important traditional ecological knowledge that has a substantial contribution for the conservation of the medicinal plants at wild. Therefore, we recommend the concerned body particularly the District Agricultural Organization to incorporate these local practices and perception to the scientific methods of conservation proposed by the Agricultural Organization for ensuring sustainable, integrated and long term management of wild medicinal plants in the study area. Moreover, active formal and/or informal local institutions shall be developed to sustain this traditional knowledge in the District.

Keywords: district lower Dir, Ethnoecology, traditional knowledge, wild medicinal plants

Introduction

Plants are among the best gifts of nature that provide the basic needs for indigenous community, especially in the developing world. They provide wood, fuel, tools, food, medicines, and fodder and grazing for livestock. In Dir Lower, traditional knowledge of plant-human relationships are as old as human history itself (Khan *et al.*, 2014, Shuaib *et al.*, 2014) ^[23, 26] and cultivation of some of these useful plants/crops is as old as the crop themselves, whose history can be traced back to the time of Queen Sheba. One of the predominant plant related traditional knowledge was those related to the wild medicinal plants; which was formally published for the first time by Poncet (1709) (Ahmad *et al.*, 2014) ^[3]. The formal documentation and publication of traditional knowledge associated with the wild medicinal plants was also continued even when the next Ethiopian governor (Emperor Mentwab) came to power. One of the remarkable travellers during the time of Emperor Mentwab was that of the Scottish James Bruce who made his travel to find the source of the Blue Nile (Thomas *et al.*, 2014) ^[9]. In his travel he collected about 160 plant species, some of which were with medicinal qualities and described together with their associated traditional medicinal knowledge. Currently, the dimension of ethnobotany got wider and encompasses number of botanical aspects of ethnoscientific studies as a result of the differences in the dimension of the knowledge of the traditional people towards their environment. Some of the botanical aspects of ethnoscientific studies are ethnomedicine, ethnotaxonomy, Ethnoecology, ethnopharmacology among many others (Ali *et*

al., 2014) ^[2]. Ethnoecology as one dimension of ethnoscientific discipline deals with the relationship of living beings (including humans) with one another and with their environment (Nazarea, 1999) ^[23]. These relationships can be social, economic, symbolic, religious, commercial and/or artistic (Bassols & Toledo 2005). Ethnoecology emphasizes on documenting traditional ecological knowledge, which may include the culture and beliefs that has been handed down through generations by cultural transmission (Haenn, 1999) ^[17]. Ethnoecological knowledge can be applied in long term management and conservation strategies of biodiversity in general (Junior and Sato, 2005) ^[24] and wild medicinal plants in particular (Aumeeruddy and Ji, 2003; Ghimire *et al.*, 2004) ^[16]. This research was initiated to show the existing of the indigenous people of District Dir Lower that has important contribution in the conservation of wild medicinal plants before it is too late to capture them.

2. Methods and Materials

The research was carried out in District, Dir Lower Khyber Pakhtun Khwa Pakistan. The District is approximately 96, 680 hectare in size. District Lower Dir are located in Khyber Pakhtunkhwa Pakistan, which the most important districts, both culturally and historically. Into two separate districts Dir was divided (Upper Dir and Lower Dir). These district came into existence. Lower Dir lies among 71° 31' to 72° 14' east longitudes and 34° 37' to 35° 07' north latitudes. From 600 m to 3000 m the altitude of the district areas. To the east it is bordered by Swat and

upper Dir, Malakand to the south and Upper Dir to the north, to the west Afghanistan and Bajaur. From northeast to southwest. The winter in the district is frosty and intolerable. In these district temperate and pleasant summer. Due to western disturbances the precipitation in the winter take place and compared to the summer the rainfall in the winter is more. With sparse forests Hill tops are covered, Pine, *Acacia*, *Ailanthus*, Oak, *Eucalyptus*, and *Olea* in these forests are the dominant trees (Shuaib *et al.*, 2014) [28].

2.1 Ethnobotanical information

Informants are representatives of the local/indigenous people of the District who can provide the ethnomedicinal information; and accordingly those informants which age ranges from 19 to 86 were chosen both randomly and systematically. Demographic figure for the population of the District whose age ranges between 18 and 85 years were in the ratio of 66: 34 for men and women. Keeping the same proportion, total of 105 informants (36 females and 69 males) were selected from 15 villages (per village 7). 60 of the total informants (per village 4) were selected randomly. By

tossing a coin and using him/her as informant every time head of the coin was up if he/she was volunteered to contribute some of them were chosen. In the selected areas some are chosen accidentally through the houses. The other 45 informants (per village 3) were local experts that were selected systematically based on references from the local people, development agents and local authorities at each research sites. Their socio-demography of informants is summarized in the following table (Table1).

2.2 Semi-structured interview

Semi-structured checklist was prepared in advance to ensure informant consensus about the traditional ecological knowledge and medicinality of each herbals following (Cunningham, 2001). The language that has been used most frequently with the informants was Pashto (common language of the District) used with the help of interpreters who were with good knowledge of the local cultures and vegetation. The interviews were done with the native people of the District.

Table 1: Socio-demographic data of the informants used in District Dir Lower

Informant parameters		Age		Sex		Education level		Informants		Distance (Km)		Marital status		Total
		Youngsters (Age between 18-30)	Elders (Age>30)	Male	Female	Literate	Illiterate	Key	Random	Lives in less than 10 Km	Lives in more than 10 Km	Married	Unmarried	
Age	Youngsters (Age between 18-30)	31												31
	Elders (Age >30)		74											74
Sex	Male	20	49											69
	Female	11	25											36
Education level	Literate	19	23	30	12									42
	Illiterate	12	51	39	24									63
Informants	Key	4	41	36	9	12	33							45
	Random	27	33	33	27	30	30							60
Distance (Km)	Lives in less than 10 Km	5	9	6	8	8	6	6	8					14
	Lives in more than 10 Km	26	65	63	28	36	57	39	52					91
Marital status	Married	12	19	25	6	18	13	2	29	9	22			31
	Unmarried	19	55	44	30	24	50	43	31	5	69			74
Total No. of informants		31	74	69	36	42	63	45	60	14	91	31	74	105

N. B. *Numbers in each cell refers to the number of informants; and distance is measured from Timergara; Number in bold refers to the total number of informants used in the study

2.3 Plant interview

This method was used to know the medicinality of each herbals collected from each study localities (Gerique, 2006). In this method, we collected the medicinal plants from the studied field area and brought them back to the village and presented to the randomly chosen informants (Figure 2) to tell us wheather these species have medicinal qualities in the District (Alexiades, 1996). By the time when we loss the live plant species, pressed specimen was used instead for this interview.

2.4 Group discussion

In each village group discussions were used for cross-checking and verifying the information that has been gathered by plant interview and semi-structured interview following (Cotton, 1996). With the local people the discussions were made, key informants and other traditional healers some-times altogether or alone in their groups throughout field Resaerch. Brief introduction was given to the groups so as to encourage them to discuss genuinely. The places and time for discussion were arranged based on the interest of the informants.

2.5 Method of data collection on wild medicinal plants

Sampling design

Out of the 27 villages in the District, 15 of them (55. 6%) were used as sampling for data collection. By purposive sampling method the selection of the 15 was made based on the availability of key informants identified with the assistance of elders and local authorities. The informants categorized the availability of wild medicinal vegetation of the District into six general habitats types, namely *Laffa Bosoona*, *Laffa Mukke*, *Laffa Choroka*, *Laffa*

Merga, *Laffa hori edu* and *Laffa ekiri*. Within each localities were identified based on these six habitats. This procedure gave 95 total localities from which 59 sampling units were selected by taking one locality for each habitat type in each village using the draw method. The selection of localities was established on stratification of habitat type it is the best symbolic sample for capturing the Ethnomedicinal knowledge and wild medicinal plant in the District.

Table 2: Total number of wild localities and number of sampling units in District Dir Lower

S/N	Villages Name	Different categories of the wild environment in District Dir Lower														Total	
		Forest land		Wood land		Wet land		Grass land		Grazing land		Fallow land					
		TNL	CNL	TNL	CNL	TNL	CNL	TNL	CNL	TNL	CNL	TNL	CNL	TNL	CNL		
1	Malaknd	NR	NR	1	1	NR	NR	1	1	1	1	1 (BL)	----	4	3		
2	Undary	NR	NR	2	1	NR	NR	2	1	1	1	1	1	6	4		
3	Maidan	NR	NR	2	1	1	1	NR	NR	2	1	1 (BL)	----	6	3		
4	Lal Qala	NR	NR	3	1	1	1	1	1	1	1	1 (BL)	----	7	4		
5	Kumbare	1	1	2	1	NR	NR	1	1	1	1	1 (BL)	----	6	4		
6	Hyaray	1	1	2	1	NR	NR	1	1	1	1	1 (BL)	----	6	4		
7	Kupardara	NR	NR	2	1	NR	NR	1	1	1	1	1 (BL)	----	5	3		
8	Haji Abad	1	1	2	1	NR	NR	2	1	1	1	1 (BL)	----	7	4		
9	Laram	1	1	2	1	1	1	1	1	1	1	1 (BL)	----	7	5		
10	Petodara	1	1	3	1	NR	NR	NR	NR	1	1	1	1	6	4		
11	Makhay	NR	NR	3	1	1	1	NR	NR	1	1	1 (BL)	----	6	3		
12	Hiasarak	1	1	2	1	1	1	1	1	2	1	1 (BL)	----	8	5		
13	Jbagay	1	1	2	1	1	1	NR	NR	1	1	1 (BL)	----	6	4		
14	Shatay	NR	NR	2	1	NR	NR	1	1	1	1	2	1	6	4		
15	Darmal	2	1	2	1	1	1	1	1	2	1	1 (BL)	----	9	5		
Total number of localities		9	----	32	----	7	----	13	----	18	----	16	----	95	----		
Total number of sampling units		----	8	----	15	----	7	----	11	----	15	----	3	----	59	----	

Keys: CNL-Chosen number of locality, NR-not represented, BL-bare land, N. B: TNL-total number of localities,

2.6 Plant identification

Medicinal plant species which were readily identifiable were recorded in the field. For those ethnomedicinal plants which were difficult to identifying in the field were provisionally kept in plastic bag; and then were pressed and brought to the Herbarium of Department of Botany, Govt Post Graduate Collage Timargara Lower Dir Pakistan. The identifications plants was done by the Flora of Pakistan. With the help of taxonomic experts in Department of Botany, Govt Post Graduate Collage Timargara Lower Dir Pakistan All the medicinal plant species were confirmed.

2.7 Method of data analysis

Descriptive statistical method such as percentage frequency method of data analysis was employed to review some of the descriptive ethnobotanical data obtained from the interviews on reported medicinal plants. Microsoft Excel spread sheet software (Microsoft Corporation, 2010) was employed for organizing and analysing some ethnobotanical data. Inferential statistical analysis using two sample independent t-test were performed to check whether there was a significant difference among the different parameters of informants (gender, age, literacy level, informant experience, marital status and living distance from health centre) for their knowledge about the medicinality of the reported medicinal plants.

3. Result

3.1 Wild ethnomedicinal vegetation composition

A total of 112 species (Table 4) of wild ethnomedicinal plants which can be grouped in to 97 genera and 53 families were documented from the study area. From these medicinal plants, 10 (9%) of them were enedemic to Dir Lower (Table 3). Among the total collected wild medicinal species, shrubs was present in highest quantity while other plants was present in least proportion (Figure 4)




Fig 4: Groth form of wild medicinal plants in District Dir Lower

Table 3: Lists of endemic wild medicinal plants in District Lower Dir

S/N	Botanical Name	Habit	Family Name
1	<i>Acacia abyssinica</i> subsp. <i>Abyssinica</i>	Tree	Fabaceae
2	<i>Inula confertiflora</i>	Shrub/tree	Asteraceae
3	<i>Impatiens rothii</i>	Herb	Balsaminaceae
4	<i>Impatiens tinctoria</i> subsp. <i>Abyssinica</i>	Herb	Balsaminaceae
5	<i>Jasminum stans</i>	Shrub	Oleaceae
6	<i>Kalanchoe petitiiana</i>	Herb	Crassulaceae
7	<i>Laggera tomentosa</i>	Shrub	Solanaceae
8	<i>Lippia adoensis</i>	Shrub	Verbenaceae
9	<i>Otostegia integrifolia</i>	Shrub	Solanaceae
10	<i>Rhus glutinosa</i> subsp. <i>neoglutinosa</i>	Shrub	Anacardiaceae
11	<i>Solanecio gigas</i>	Shrub	Asteraceae
12	<i>Solanum marginatum</i>	Shrub	Solanaceae
13	<i>Thymus schimperi</i>	Herb	Lamiaceae
14	<i>Urtica simensis</i>	Herb	Urticaceae

Table 2: Lists of wild medicinal plants collected from District Lower

0	Scientific Name	Family	Growth form	U T	Ailment treated
1	<i>Acacia abyssinica</i> Hochst. ex Benth.	Fabaceae	T	An	Horse scabies
2	<i>Acacia albida</i> Del.	Fabaceae	T	An	Eye bruise (Bilz)
3	<i>Acacia seyal</i> Del.	Fabaceae	T	Hu	Headache (Ras mitat)
4	<i>Achyranthes aspera</i> L.	Amaranthaceae	H	Hu	Stomach trouble
				Hu	Abdominal pain in woman after birth
				Hu	RH case (Shotelay)
5	<i>Acmella caulirhiza</i> Del.	Asteraceae	H	Hu	Loose tooth
6	<i>Agave sisalana</i> Perrine ex Engel.	Agavaceae	T	An	Tick
7	<i>Ageratum houstonianum</i> Mill	Asteraceae	H	Hu	Poisoning (Merzenet)
8	<i>Ajuga integerifolia</i> Buch. Ham.	Lamiaceae	H	Hu	Stomach trouble
				Hu	Cold (Bired)
				Hu	Gout (Rihi)
				Hu	Hypertension (Dem bizat)
9	<i>Aloe macrocarpa</i> Tod.	Aloaceae	H	Hu	Intestinal parasite
				Bo	Swelling (Ebach)
10	<i>Alternanthera pungens</i> Kunth.	Amaranthaceae	H	Hu	Sudden illness (Dingetegna)
11	<i>Artemisia abyssinica</i> Schtz. Bip. ex Rich	Asteraceae	H	Hu	Whooping Cough (Tektik)
				Hu	Stomach trouble
				Hu	Eye itching (Ayenen masakek)
12	<i>Asparagus africanus</i> Lam.	Asparagaceae	S	Hu	Amobiasis (Ameba)
13	<i>Asparagus racemosus</i> Wild.	Asparagaceae	S	Hu	Amobiasis
14	<i>Asplenium monanthes</i> L.	Aspleniaceae	H	Hu	Womb itching (Mehatsenen masakek)
15	<i>Bersama abyssinica</i> Fresen.	Meliantaceae	T	An	Horse Scabies (Bech'h)
16	<i>Bidens pilosa</i> L.	Asteraceae	H	Hu	Devil sickness (Lekefet)
17	<i>Brucea antidysenterica</i> J. F. Mill.	Simaroubaceae	S	Hu	Evil eye (Buda)
				An	Colic (yehod hemem)
18	<i>Buddleja polystachya</i> Fresen.	Buddlejiaceae	T	An	Leech (Alekit)
19	<i>Calpurnia aurea</i> (Ait.) Benth.	Fabaceae	S	Hu	Scabies (Ekek)
				An	Pubic hair louse (Qemanjer)
20	<i>Capparis tomentosa</i> Lam.	Capparidaceae	CL	Hu	Wound (Kusil)
21	<i>Carissa spinarium</i> (Vahl.) Forssk. ex Endl.	Apocynaceae	S	Hu	Intestinal worms
				Hu	Evil eyes
22	<i>Centella asiatica</i> (L.) Urban.	Apiaceae	H	Hu	Bleeding
23	<i>Clausena anisata</i> (Wild.) Benth.	Rutaceae	S	Hu	Toothache
24	<i>Clematis simensis</i> Fresen.	Ranunculaceae	LI	Bo	Wound
				Hu	Evil eye
				Hu	Wart (Kintarot)
25	<i>Clerodendrum myricoides</i> (Hochst) Vatke	Lamiaceae	S	Hu	Diarrhae
26	<i>Colocasia esculenta</i> (L.) Schott	Araceae	H	Hu	Swelling
27	<i>Croton macrostachyus</i> Del.	Euphorbiaceae	T	Hu	Febril illness (Megagna)
				Hu	Tinea nigra (Kuakucha)

28	<i>Cucumis dipsaceus</i> Ehrenb.	Cucurbitaceae	CL	Hu	Depression (Eje seb)
29	<i>Cucumis ficifolius</i> A. Rich.	Cucurbitaceae	CL	Hu	Abdominal pain (Kuretet)
30	<i>Cyathula cylindrica</i> Moq.	Amaranthaceae	H	Hu	Stomachache (Yehod hemem)
31	<i>Cyphostemma adenocaula</i> (Steud. ex A. Rich.) Descouings ex Wild & Drummond	Vitaceae	CL	An	Blackleg
				Bo	Swelling
				Hu	Snake bite
32	<i>Datura stramonium</i> L.	Solanaceae	H	Hu	For Intellegency (Letimret)
33	<i>Dodonaea angustifolia</i> L. f.	Sapindaceae	S	An	Wound
34	<i>Dombeya torrida</i> (J. F. Gmel) Bamps	Sterculiaceae	T	Hu	Antidot for snake bites
35	<i>Dregea schimperi</i> (Decne.) Bullock	Asclepiadaceae	LI	Hu	Eczema (Chiffea)
36	<i>Ekebergia capensis</i> Sparrm.	Meliaceae	T	Hu	Syphilis (Kitign)
37	<i>Eleusine floccifolia</i> Forssk.	Poaceae	H	Hu	Snake bit
				Hu	Poisoning
38	<i>Embelia schimperi</i> Vatke	Myrsinaceae	S	Hu	Tape worm (Kosso)
39	<i>Erica arborea</i>	Ericaceae	S	An	Eye disease
40	<i>Euclea racemosa</i> subsp. <i>schimperi</i>	Ebenaceae	S	Hu	Tonsillitis (Entil siwored)
41	<i>Euphorbia abyssinica</i> J. F. Gmel.	Euphorbiaceae	T	Hu	Haemorrhage
42	<i>Euphorbia ampliphylla</i>	Euphorbiaceae	T	Hu	Haemorrhage
43	<i>Ferula communis</i> L.	Apiaceae	H	Hu	Cough
44	<i>Ficus sur</i> Forssk.	Moraceae	T	Hu	Wart on hand(Kintarot)
				An	Swelling
45	<i>Foeniculum vulgare</i> Mill.	Apiaceae	H	Hu	Urinary Retention (Shinet leklelelew)
				Hu	Stomach trouble
46	<i>Fuerstia africana</i> Th. Fries	Lamiaceae	H	Hu	General malaise (Mich)
				An	Cattle eye disease
47	<i>Gamphocarpus abyssinicus</i> Decne.	Asclepiadaceae	H	An	Blackleg (Aba gorba)
48	<i>Grewia ferruginea</i> Hochst ex . A . Rich.	Tiliaceae	S	Hu	Taeniasis (Kosso)
49	<i>Guizotia scabra</i> (Vis) Chiov.	Asteraceae	H	Hu	Epilopsy (Yemitel beshita)
50	<i>Heteromorpha trifoliata</i> (Wendel) Eckl. & Zeyh.	Apiaceae	S	Hu	Warding of Sorcery Stealing (Selabi)
51	<i>Hygrophila schulli</i> (Hamilt.) M. R. & S. M. Almeida	Acanthaceae	H	Bo	Poisoning
52	<i>Hypericum quartianum</i> A. Rich.	Hypericaceae	S	Hu	Jaundice (Yewof beshita)
53	<i>Hypericum revolutum</i> Vahl	Hypericaceae	S	An	Eye disease
54	<i>Impatiens rothii</i> Hook. f.	Balsaminaceae	H	Hu	Wounds on hand
55	<i>Impatiens tinctoria</i> A. Rich. Subsp. <i>abyssinica</i> (Hook. f.) Grey-Wilson	Balsaminaceae	S	Hu	Wound on palm
56	<i>Inula confertiflora</i> A. Rich.	Asteraceae	S	An	Eye disease
				Bo	Rabies (Yehebid wusha beshita)
57	<i>Jasminum grandiflorum</i> L.	Oleaceae	S	Hu	Evil eye
				Hu	Toothache (Yeters himem)
58	<i>Juniperus procera</i> Endle	Cupressaceae	T	Hu	Demon possesion (Ganen)
59	<i>Kalanchoe petitiiana</i> A. Rich.	Crassulaceae	H	Bo	Swelling
60	<i>Lagenaria siceraria</i> (Molina) Standl.	Cucurbitaceae	H	Hu	Impotency (Sinfet wosib)
61	<i>Laggera tomentosa</i> (Sch. Bip. ex A. Rich.) Oliv. & Hiern	Asteraceae	H	Hu	Flu (Gunfan)
62	<i>Leonotis raineriana</i> Vis.	Lamiaceae	S	An	Leech
				Hu	General malaise (Mich)
63	<i>Leucas martinicensis</i> (Jacq.) R. Br.	Lamiaceae	S	Hu	General malaise (Mich)
64	<i>Lippia adoensis</i> Hochst. ex Walp.	Verbenaceae	S	Hu	Stomach pain (Cheguara)
65	<i>Maesa lanceolata</i> Forssk.	Myrsinaceae	S	Bo	Swelling
66	<i>Malva venticillata</i> L.	Malvaceae	H	An	Swelling
67	<i>Myrica salicifolia</i> A. Rich.	Myricaceae	T	Hu	Ascariasis
68	<i>Myrsine africana</i> L.	Myrsinaceae	S	Hu	Taeniasis
				An	Worms in donkey
69	<i>Ocimum gratissimum</i> L.	Lamiaceae	H	Hu	General malaise
70	<i>Ocimum lamifolium</i> Hochst. ex Benth.	Lamiaceae	S	Hu	General malaise
71	<i>Olea europaea</i> L. subsp. <i>cuspidata</i> (Wall. ex G. Don) Cif.	Oleaceae	T	Hu	Snake bite
72	<i>Olinia rochetiana</i> A. Juss.	Oliniaceae	S	Hu	Snake bit
				Hu	Toothache
73	<i>Osyris quadripartita</i> Decn.	Santalaceae	S	Hu	Eczema
74	<i>Otostegia integrifolia</i> Benth.	Lamiaceae	S	Hu	Fibril illness (Megagna)
75	<i>Pavetta abyssinica</i> Fresen.	Rubiaceae	S	Hu	Poison
				An	Animal diarrhoea

76	<i>Pentas schimperiana</i> (A. Rich.) Vatke	Rubiaceae	S	An	Eye disease
77	<i>Phytolacca dodecandra</i> L' Herit	Phytolaccaceae	S	An	BECH'E'H
				Hu	Wart on hand
78	<i>Plantago lanceolata</i> L.	Plantaginaceae	H	An	Intestinal parasite
79	<i>Plantago major</i> L.	Plantaginaceae	H	Hu	Poisoning
				Hu	Haemorrhoides
80	<i>Premna schimper</i> Engl	Lamiaceae	S	Hu	Eye disease
81	<i>Protea gaguedi</i> J. F. Gmel.	Proteaceae	S	An	Animal jaundice
82	<i>Prunus africana</i> (Hook. f.) Kalms	Rosaceae	T	Hu	Swelling
				Hu	Sudden illness (Dingetegna)
				An	Blackleg
				An	Anthrax (Abasenga)
83	<i>Pterolobium stellatum</i> (Forssk.) Brenan	Fabaceae	S	Hu	Rhumantic pain (Kurtimat)
84	<i>Rhus glutinosa</i>	Anacardiaceae	S	Hu	Epistaxis (Neser)
85	<i>Rhus retinorrhoea</i>	Anacardiaceae	S	An	Anthrax (Abasenga)
86	<i>Rhus vulgaris</i> Meikle	Anacardiaceae	S	An	Diarrhoea
87	<i>Ricinus communis</i> L.	Euphorbiaceae	T	Hu	Dandruff (Forofor)
88	<i>Rosa abyssinica</i> Lindley	Rosaceae	S	An	Invoking spirit (Aganent)
89	<i>Rubia cordifolia</i> L.	Rubiaceae	H	Hu	Wound
				Hu	Cough
				Hu	Cough
				An	Cataract (Bemora yete-shefene ayen)
90	<i>Rubus steudneri</i> Schweinf.	Rosaceae	H	Hu	Stabbing pain (Wugat)
				Hu	Cough
91	<i>Rumex abyssinicus</i> Jacq.	Polygonaceae	H	Hu	Eye bruise
				An	Blackleg
				An	Scabies (Ekek)
92	<i>Rumex nepalensis</i> Spreng.	Polygonaceae	H	An	Colic (Yehod himem)
				An	Blackleg
				Hu	Stomach pain (Cheguara)
				Hu	Stabbing pain (Wugat)
				B	Urinary retention
93	<i>Rumex nervesus</i> Vahl	Polygonaceae	S	Hu	Delay in drying circumcision
94	<i>Salix mucronata</i>	Salicaceae	T	Hu	MIKEGNA-SHEREGNA
95	<i>Salvia nilotica</i> Jacq.	Lamiaceae	H	Hu	Wound
96	<i>Sida schimperiana</i> Hochst. ex A. Rich.	Malvaceae	H	An	Rabies
				An	Preventing bitch birth
97	<i>Snowdenia polystachya</i> (Fresen.) Pig.	Poaceae	H	Hu	Scabies (Ekek)
98	<i>Solanecio gigas</i> (Vatke.) C. Jeffrey	Asteraceae	S	Bo	Retained placenta
99	<i>Solanum anguivi</i> Lam.	Solanaceae	S	Hu	Intelligence
				Hu	Dandruff
				An	Rabies
100	<i>Solanum incanum</i> L.	Solanaceae	S	An	Tick bite
				An	Horse Scabies
				Hu	Wounds
101	<i>Solanum marginatum</i> Linn. f.	Solanaceae	S	Hu	Long stay menstruation
102	<i>Stephania abyssinica</i> (Dillon ex A. Rich.) Walp.	Menispermaceae	LI	An	Rabies
				An	Blackleg
				Hu	Unwanted pregnancy
				Hu	Wound
				Hu	Swelling
				Hu	Sudden illness
103	<i>Tagetes minuta</i> L.	Asteraceae	S	An	KINKIN
104	<i>Thunbergia alata</i> Sims.	Acanthaceae	CL	Hu	Cough
105	<i>Thymus schimper</i> Ronniger	Lamiaceae	S	Hu	Hypertension
106	<i>Urtica simensis</i> Steudel	Urticaceae	H	Hu	Gonorrhoea (Chebit)
107	<i>Verbascum sinaiticum</i> Benth.	Scrophulariaceae	H	Hu	Nightmare
				An	Blackleg
108	<i>Verbena officinalis</i> L.	Verbenaceae	H	Hu	Cough
				Hu	Tonsillitis (Entil siwored)
109	<i>Vernonia amygdalina</i> Del.	Asteraceae	T	Hu	Warding off sorcery stealing

				Hu	Malaria
				Hu	Abdominal pain
110	<i>Withania somnifera</i> (L.) Dunal.	Solanaceae	S	Hu	Daemon possesion
111	<i>Xanthium strumarium</i> L.	Asteraceae	S	An	Leech
112	<i>Zehneria scabra</i> L.	Cucurbitaceae	LI	Hu	Deformed lips (Megagna)
				Hu	Urinary retention

3.2 Statistical test on the reported number of medicinal plant

The questionnaire respondents represented a diverse array of people including farmers, women, literate, illiterate, youngsters, elders, married and unmarried. Among the 105 informants, 69 (65.7%) were male and 36 (34.3%) were females. The largest proportions of the respondents were elders (70.5%) above 30 years old (Table 5). Most respondents were not able to write and read (60%) whereas about 40% of the respondents were joined at

least formal grade one school and able to write or read. Among the respondents 86.7% of them dwell far away (> 5 Km) from the centre of health centres and/Timeragar DHQ center whereas only few (13.3%) of them were living near (< 5 Km) to the Timeragar DHQ center. Inferential statistical test of significance on average number of reported wild medicinal plants by the different groups of informants in District Dir Lower is shown below (Table 5)

Table 3: Average number of wild medicinal plants (AWMP) reported by different groups of informants

Parameters	Informant groups	Number of informants (n)	Percentage (%)	Total number of citations (N)	AWMP \pm SD	t-value	P-value*
Gender	Male	69	65.7%	351	34.07 \pm 20.47	-1.445	0.149
	Female	36	34.3%	191	36.74 \pm 20.56		
Age	Youngsters	31	29.5%	209	36.11 \pm 19.87	-0.985	0.325
	Elders	74	70.5%	333	34.33 \pm 20.92		
Literacy level	Illiterate	63	60%	181	35.03 \pm 21.00	-0.012	0.991
	Literate	42	40%	361	35.01 \pm 20.31		
Informant category	Key	45	42.9%	223	35.68 \pm 20.42	0.633	0.527
	General	60	57.1%	319	34.55 \pm 20.62		
Marriage	Married	31	29.5%	516	34.83 \pm 20.53	-0.925	0.355
	Unmarried	74	70.5%	26	38.65 \pm 20.47		
Distance from health centre	Close to the health centre	14	13.3%	42	44.50 \pm 18.29	-3.142	0.002*
	Far from the health centre	91	86.7%	500	34.22 \pm 20.52		

4. Discussion

Our results with regard to wild medicinal plant composition suggest that the District was once primarily a typical dry Afromontane Forest ecosystem of Dir Lower. This is due to the presence of remnant characteristic species for the vegetation type of dry evergreen Afromontane Forests (Friis *et al.*, 2011) [14]. These species include *Croton macrostachyus*, *Juniperus procera*, *Olea europaea* subsp. *cuspidata* locally called *Bekanis*, *Gatira* and *Ejersa* respectively; and are still retained in the existing landscape of the District as medicinal herbals. The local people uses a taxonomically diverse group of wild medicinal plants, about 112 species in 97 genera and 53 families. The availability of diverse medicinal plant species at wild were also reported from all corner of Dir Lower (Shinwari *et al.*, 2011) [27]. This attributed the fact that wild habitates are the main storehouse of medicinally useful plants. Some of these medicinal plants were recorded to be economically important plants used for many other purposes in the District. For example, *Juniperus procera*, *Acacia albida*, *Croton macrostachyus*, *Olea europaea* subsp. *cuspidata*, and *Prunus africana* were some of the medicinal plants in the District with multiple purposes other than their medicinal values (Khan *et al.*, 2014) [21]. Some of the medicinal plants identified in this study were reported elsewhere to have other use values other than their therapeutic quality. For example, *Cordia africana*, *Croton macrostachyus*, *Juniperus procera*, *Prunus africana*, *Olea europaea*, *Ekibergia capensis* were reported for the purpose of timber in different areas of Dir Lower (Khan *et al.*, 2013) [21].

Similarly other medicinal species such as *Acacia abyssinica*, *Acacia albida*, *Acacia seyal* were also reported elsewhere for home garden agro-forestry purposes such as fencing and shading whereas *Euphorbia ampliphyla*, *Euphorbia abyssinica*, *Croton macrostachyus*, *Vernonia amydalina* were recorded for their purpose of beehive making and/or bee forage (Hussain *et al.*, 2014)

The finding of this study showed that shrubby herbals were the most dominant form of wild medicinal plants in the District followed by herbaceous forms. Similar findings were noted elsewhere in Lower Dir (Ahmad *et al.*, 2013) this may be linked with the custom of the local people to use plants that are available almost all the time. In line with this fact, suggested that the knowledge in relation to medicinal plants directly emanates/originates from the type of the plants they are surrounded by. In this regard, shrubby herbals are the most available form of herbals in almost all year as they are tolerant to seasonal variation (Albuquerque, 2006) [5] and might have had high chance of being chosen by the local people of the study area. On the contrary the ecological natures of herbaceous medicines are normally annuals and subjected to be influenced by small scale environmental variations than shrubs (Qasim *et al.*, 2013) [26]. Moreover, apart from seasonal variation, grazing intensity in the study area might have contributed for the lesser number of herbaceous medicinal plants than shrubs (Khan *et al.*, 2014) [21]. This effect of grazing on herbaceous medicinal plants were also noted elsewhere (Adnan and Holscher, 2010) [1]. The rather fewer

contribution of trees for therapeutic purpose in the District may be linked with the less abundance of tree species resulted from the previous over exploitation and habitat modification history of trees mainly for the purposes other than their medicinal values (. The plant families such as Asteraceae (11 species, 9.82%), Lamiaceae (11 species, 9.82%), Fabaceae (5 species, 4.46%), Solanaceae (5 species, 4.46%), Apiaceae (4 species, 3.57%), Cucurbitaceae (4 species, 3.57%) and Euphorbiaceae (4 species, 3.57%) are found. This deduces the high contribution of these plant families to the majority of medicinal flora of the country. The relatively high contribution of these families other than other families may originate from their high species richness in the Flora of Dir Lower (Barkatullah *et al.*, 2015) ^[10] have found strong positive correlation ($r=0.88$) between the overall species richness of vegetation and the associated ethnomedicinal plant species richness in Pakistan. Moreover, critical observation on the joint study of vegetation and ethnomedicinal plant diversity showed that the species richness of both the general vegetation and medicinal flora follow the same pattern and seem to be directly related. In line with this concluded the presence of greater concentration of medicinal plant diversity at the areas where there is higher concentration of biological and cultural diversity. The finding of this study showed that about 10% of the collected medicinal plants are endemic to Dir Lower, which follows almost the same proportion of endemism for the Flora of Paksiatn and Dir Lower (Ahmad *et al.*, 2105) ^[3].

5. Conclusion and recommendation

This study documented 112 wild medicinal plants that can be grouped into 97 genera and 53 families. This study also found out the presence of traditional perspectives and cultural beliefs which would maintain the ecology of medicinal plant species. Thus, integrating the ethnoecological perspectives of the local/indigenous people would be helpful for better ecosystem management in general and wild medicinal plants in particular. Moreover, active formal and/or informal local institutions shall be developed to sustain this traditional knowledge in the District.

6. Conflict of interests

The authors have not declared any conflict of interests.

7. Acknowledgement

We are grateful to the Department of Botany, Govt Post Graduate Collage Timargara Lower Dir Pakistan for his moraly support. We also thank the Local Hakeem for identification of herbals. The local informants are highly appreciated for their contribution in the field work.

8. References

1. Adnan M, Hölscher D. Medicinal plant abundance in degraded and reforested sites in Northwest Pakistan. *Mountain Research and Development*. 2010; 30(1):25-33.
2. Ahmad I, Ibrar M, Ali N. Ethnobotanical study of Tehsil Kabal, Swat District, KPK, Pakistan. *Journal of Botany*, 2011.
3. Ahmad L, Semotiuk A, Zafar M, Ahmad M, Sultana S, Liu QR, Yaseen G. Ethnopharmacological documentation of medicinal plants used for hypertension among the local

- communities of DIR Lower, Pakistan. *Journal of ethnopharmacology*. 2015; 175:138-146.
4. Ahmad M, Sultana S, Fazl-i-Hadi S, Ben Hadda T, Rashid S, Zafar M, Yaseen G. An Ethnobotanical study of Medicinal Plants in high mountainous region of Chail valley (District Swat-Pakistan). *Journal of ethnobiology and ethnomedicine*. 2014; 10(1):36.
5. Albuquerque UP, Lucena RF, Monteiro JM, Florentino AT, Cecília de Fátima CBR. Evaluating two quantitative ethnobotanical techniques. *Ethnobotany Research and Applications*. 2006; 4:051-060.
6. Alexiades MN. Collecting ethnobotanical data: an introduction to basic concepts and techniques. *Advances in Economic Botany*. 1996; 10:53-94.
7. Ali M, Begum R, Ali K, Nisar M, Khan W, Hassan N, Khan N. Ethnobotanical survey of wild medicinal plants of Tirat Valley District, Swat, Khyber-Pakhtunkhwa. *J Biodiversity Environmental Sci*. 2017; 11:91-101.
8. Aumeeruddy-Thomas Y, Pei SJ. *Applied ethnobotany: case-studies from the Himalayan region*. WWF, 2003.
9. Aumeeruddy-Thomas Y, Shinwari ZK, Abdullah A, Khan AA. Ethnobotany and the management of fodder and fuel wood at Ayubia National Park, North West Frontier Province, Pakistan. *People and Plants working paper*, 2004, 13.
10. Barkatullah IM, Rauf A, Hadda TB, Mubarak MS, Patel S. Quantitative ethnobotanical survey of medicinal flora thriving in Malkand Pass Hills, Khyber PakhtunKhwa Pakistan. *Journal of Ethnopharmacology*. 2015; 169:335-346.
11. Barrera-Bassols N, Toledo VM. Ethnoecology of the Yucatec Maya: symbolism, knowledge and management of natural resources. *Journal of Latin American Geography*. 2005; 4(1):9-41.
12. Beck K, Beedle M, Van Bennekum A, Cockburn A, Cunningham W, Fowler M, Kern J. *Manifesto for agile software development*, 2001.
13. Cotton CM, Wilkie P. *Ethnobotany: principles and applications* (No. Sirsi) i9780471955375). Chichester: John Wiley & Sons, 1996.
14. Friis EM, Crane PR, Pedersen KR. *Early flowers and angiosperm evolution*. Cambridge University Press, 2011.
15. Gerique A. An introduction to ethnoecology and ethnobotany: Theory and methods. Integrative assessment and planning methods for sustainable agroforestry in humid and semiarid regions. *Advanced Scientific Training*. Loja, 2006.
16. Ghimire SK, McKey D, Aumeeruddy-Thomas Y. Conservation of Himalayan medicinal plants: Harvesting patterns and ecology of two threatened species, *Nardostachys grandiflora* DC and *Neopicrorhiza scrophulariiflora* (Pennell) Hong. *Biological Conservation*. 2005; 124(4):463-475.
17. Haenn N. The power of environmental knowledge: Ethnoecology and environmental conflicts in Mexican conservation. *Human Ecology*. 1999; 27(3):477-491.
18. Hussain I, Ullah R, Khurram M, Ullah N, Baseer A, Khan FA, Khan J. Heavy metals and inorganic constituents in medicinal plants of selected districts of Khyber

- Pakhtoonkhwa, Pakistan. African Journal of Biotechnology. 2011; 10(42):8517-8522.
19. Khan, AA, Ali F, Ihsan M, Hayat K, Nabi G. Ethnobotanical study of the medicinal plants of Tehsil Charbagh, district Swat, Khyber Pakhtunkhwa, Pakistan. Am.-Eurasian J. Agric. Environ. Sci. 2015; 15:1464-1474.
 20. Khan I, Abdel Salam NM, Fouad H, Tariq A, Ullah R, Adnan M. Application of ethnobotanical indices on the use of traditional medicines against common diseases. Evidence-Based Complementary and Alternative Medicine, 2014.
 21. Khan N, Abbasi AM, Dastagir G, Nazir A, Shah GM, Shah MM, Shah MH. Ethnobotanical and antimicrobial study of some selected medicinal plants used in Khyber Pakhtunkhwa (KPK) as a potential source to cure infectious diseases. BMC complementary and alternative medicine. 2014; 14(1):122.
 22. Khan RU, Mehmood S, Jaffar SUKF. Ethnobotanical study of food value flora of district Bannu Khyber Pakhtunkhwa, Pakistan. Journal of Medicinal Plants. 2013; 1:4.
 23. Nazarea VD. (Ed.). Ethnoecology: situated knowledge/located lives. University of Arizona Press, 1999.
 24. Pedroso-Júnior NN, Sato M. Ethnoecology and conservation in protected natural areas: incorporating local knowledge in Superagui National Park management. Brazilian Journal of Biology. 2005; 65(1):117-127.
 25. Pohle P, Gerique A. Traditional ecological knowledge and biodiversity management in the Andes of southern Ecuador. Geographica Helvetica. 2006; 61(4):275-285.
 26. Qasim M, Khalid M, Sayyed A, Din I, Hayat K, Jan SA. Phytochemical potentials and medicinal uses of twenty-four selected medicinal plants from Swabi, Pakistan. Journal of Rural Development and Agriculture. 2016; 1(1):49-58.
 27. Shinwari SHEHLA, Qureshi R, Baydoun E. Ethnobotanical study of Kohat Pass (Pakistan). Pakistan Journal of Botany. 2011; 43(SD):135-139.
 28. Shuaib M, Khan I, Sharifullah RK, Hashmatullah SM, Naz R. Ethnobotanical studies of spring flora of Dir Lower, Khyber Pakhtunkhwa, Pakistan. Pak. J. Weed Sci. Res. 2014; 20(1):37-49.