



International Journal of Agriculture and Nutrition

www.agriculturejournal.net

Online ISSN: 2664-6072; Print ISSN: 2664-6064

Received: 02-12-2019; Accepted: 04-01-2020; Published: 13-01-2020

Volume 2; Issue 1; 2020; Page No. 08-12

Identifying bread wheat lines for high zinc, iron and low phytate concentration

Habtamu Gebremichael¹, Melaku Tafese^{2*}, Habtemariam Zegeye³, Abraha Gebregiorgis⁴, Dugasa Gerenfess⁵, Neima Demsis⁶, Cherinet Kassahu⁷, Bilatu Agza⁸

^{1, 2, 3, 4, 6, 7, 8} Kulumsa Agricultural Research Center, 8 Ethiopian Institute of Agricultural Research Head Quarter, EIAR, P.O. Box 2003, Addis Ababa, Ethiopia

DOI: <https://doi.org/10.33545/26646064.2020.v2.i1a.24>

Abstract

Micronutrient malnutrition, “hidden hunger” affects about two billion people worldwide. It is responsible for escalated morbidity and mortality rates, reduced children’s cognitive ability, diminished labor power productivity and higher chronic diseases prevalence. Zinc deficiency causes impairment of physical growth, immune system and learning ability, increased risk of infections, DNA damage and cancer development. Meanwhile, iron deficiency results in tissue hypoxia and heart failure, maternal anemia, stunted and unhealthy babies (born from iron deficient mothers), children with poor attention spans, impaired fine motor skills, less capacity for memory and less work productivity. Nutrition plays a key role in improving quality of agricultural produce and well-being and productivity of the society, hence crop breeding using genetic bio fortification is best option as it reaches every house hold using the seed for cultivation specially bread wheat seed is the most common cultivated cereal in different parts of the country, thus iron and zinc bio fortified wheat will reach every household within the baked bread, thus harvest plus wheat samples provided by Kulumsa agricultural research center, wheat breeding program and the appropriate level of the micro nutrients iron (Fe) and zinc (Zn) determined by Atomic Absorption Spectrophotometer (AAS) and with the anti-nutritional factors (phytate and tannin) were also assessed using UV-Vis spectrophotometer and with respect to upper limit Recommended Daily Allowance (RDA) values per day without any possible effects of overdose intake 8HPAN121(Fe,16.94 mg/ 100g; Zn,1.218 mg/100g), 8HPAN19 (Fe,16.26 mg/ 100g; Zn,1.634 mg/100g), 7HPAN31(Fe,15.64 mg/100 g; Zn,1.97 mg/100 g) and 7HPYT44 (Fe, 17.36 mg/ 100g; Zn, 2.29 mg/ 100 g) of the harvest plus bio fortified wheat had shown promising results since the iron contents had shown to be safer to be consumed twice per day per 100 gram bread considering other sources of iron in day to day food consumed, relatively better zinc content and lower phytate contents consequently be a better candidate for parental cross breeding, adaptability and stability studies in different agro ecological zones and seasons.

Keywords: bread wheat, bio fortified wheat, harvest plus, iron, micronutrient, phytate, RDA, spectrophotometer, tannin, zinc

Introduction

Micronutrient malnutrition, “hidden hunger” affects about two billion people worldwide. It is responsible for escalated morbidity and mortality rates, reduced children’s cognitive ability, diminished labor power productivity and higher chronic diseases prevalence. Zinc and iron deficiency causes mortality of 500,000 under 5 year children annually (Velu *et al.*, 2012) ^[12]. Among the micronutrients, zinc deficiency causes impairment of physical growth, immune system and learning ability, increased risk of infections, DNA damage and cancer development (Cakmak, 2008; Gibson, 2006; Welch and Graham, 2002). Meanwhile, iron deficiency results in tissue hypoxia and heart failure, maternal anemia, stunted and unhealthy babies (born from iron deficient mothers), children with poor attention spans, impaired fine motor skills, less capacity for memory and less work productivity (Sramkova *et al.*, 2009). Phytates (myo-inositol hexaphosphate) reduces bioavailability of minerals such as Calcium, Iron, Zinc and Magnesium. Most of the inorganic phosphorus (Pi) present in mature cereal seeds (40–80%) is stored as phytate (Sramkova *et al.*, 2009). Genetic bio fortification assures a sustainable, cost-effective alternative to conventional fortification (Welch and Graham, 2002). Thus the aim of the study was to identify and select CIMMYT imported

bread wheat lines for high zinc and Iron concentration and lower phytate content for the minerals bioavailability and to select nutrition wise the appropriate genotypes for bread wheat cross breeding program at Kulumsa Agricultural Research Center.

General Objective

1. To identify CIMMYT imported bread wheat lines with low phytate and high minerals concentration.

Specific Objectives

1. To analyze the mineral concentration (iron and zinc) of the bread wheat lines.
2. To analyze the phytate and tannin content of the bread wheat lines.
3. To identify the bread wheat lines with higher amount of Iron and Zinc with lower phytate content for the parental crossing in the bread wheat breeding program.

Materials and Methods

Study Area

CIMMYT imported Harvest Plus bread wheat lines having better yield and resistant to diseases were collected from Kulumsa Agricultural Research Center, bread wheat breeding program.

The analysis was carried out at Holetta Agricultural & Nutritional Research Laboratory (iron and zinc analysis), Debrezeit Agricultural Research Center (Dry ashing and sample extraction), Kulumsa Food Science and Nutrition laboratory (Sample milling) and Melkassa Agricultural Chemistry Laboratory (Phytate and tannin analysis). Completely Randomized design (CRD) was used.

Sample preparation for iron and zinc analysis

About 1.0g of ground Harvest Plus bread wheat sample was put in aluminium dish over night at 105°C in an oven and cooled in a desiccator. Then exactly 0.5 g of dried sample was put into a porcelain crucible and then put it in a muffle furnace, the temperature adjusted to 450°C, at this temperature ashed for 4 hours, and then allowed to cool inside the closed furnace overnight. The ashed material was transferred into 200 ml Erlenmeyer flask with 20 ml 20% HNO₃. Then the acid treated sample was heated on hot plate to boiling and maintained at this temperature for 30 minutes with periodic stirring by glass rod, and allowed cool. The sample was then filtered through a no. 1 filter paper into a 100 ml volumetric flask. The contents are further washed until 90 ml of the filtrate collected, and bring to volume with distilled water. The solution was then used for the determination of iron and zinc in the Harvest Plus wheat samples. The standard curve was done and the absorbance of iron (Fe) and zinc (Zn) were measured with Flame Atomic Absorption Spectrophotometer (AAS) (Perkin-Elmer 560, Norwalk, CT) at 248.3 nm and 213.9 nm respectively. (Menaleshoa and Sorsa, 2010) [8].

$$\text{Fe and Zn (ppm)} = \frac{(a-b) * V * mcf}{s}$$

Where a = Concentration of (Fe and Zn) in plant digest from AAS reading (ppm).

b = Concentration of (Fe and Zn) in blank digest from AAS reading (ppm).

V = Total volume of digest (100 ml).

s = Weight of sample (g).

mcf = Moisture correction factor.

Determination of phytate content

About 0.1000 g of fresh samples were extracted with 10 ml 2.4% HCl in a mechanical shaker (Eberbach) for 1 hour at an ambient temperature and centrifuged at 3000 rpm for 30 min. The clear supernatant was used for phytate estimation. A 2 ml of Wade reagent (containing 0.03% solution of FeCl₃.6H₂O and 0.3% of sulfosalicylic acid in water) was added to 3 ml of the sample solution (supernatant) and the mixture was mixed on a Vortex (Maxi Maxi II) for 5 s. The absorbance of the sample solutions were measured at 500 nm using UV-vis spectrophotometer (Beckman DU-64-spectrophotometer, USA). A series of standard solution were prepared containing 0, 5, 10, 20 and 40 µg/ml of phytic acid (analytical grade sodium phytate) in 0.2 N HCl. A 3 ml of standard was added into 15 ml of centrifuge tubes with 3 ml of water which were used as a blank. A 1 ml of the Wade

reagent was added to each test tube and the solution was mixed on a Vortex mixer for 5 s. The mixtures were centrifuged for 10 min and the absorbances of the solutions (both the sample and standard) were measured at 500 nm by using deionized water as a blank. A standard curve was made from absorbance vs. concentration. (Bulbula and Urga, 2018) [1].

Determination of Condensed tannin

Tannin content was determined by the method of Burns (Burns, 1971). About 2.0000 g of chickpea flour was weighed in a screw cap test tube. The chickpea flour was extracted with 10 ml of 1% HCl in methanol for 24 h at room temperature with mechanical shaking. After 24 h shaking, the solution was centrifuged at 1,000 rpm for 5 min. A 1 ml of supernatant was taken and mixed with 5 ml of vanillin-HCl reagent (prepared by combining equal volume of 8% concentrated HCl in methanol and 4% Vanillin in methanol). D-catechin was used as standard for condensed tannin determination. A 40 mg of D-catechin was weighed and dissolved in 1,000 ml of 1% HCl in methanol, which was used as stock solution. A 0, 0.2, 0.4, 0.6, 0.8 and 1 ml of stock solution was taken in test tube and the volume of each test tube was adjusted to 1 ml with 1% HCl in methanol. A 5 ml of vanillin-HCl reagent was added into each test tube. After 20 min, the absorbance of sample solutions and the standard solution were measured at 500 nm by using water to zero the spectrophotometer (Bulbula and Urga, 2018) [1].

Statistical analysis

The mean and standard deviation (SD) of means were calculated. The data were analyzed by one-way analysis of variance (ANOVA) and mean comparison was done with Least Significant Difference and correlation with person correlation coefficient using Minitab version 17 software. A level of p < 0.05 was used to indicate significant differences among the samples.

Results and Discussion

As shown in table 1 below iron Recommended Dietary Allowance (RDA) for men is 8 mg/ day and 18 mg /day for women of 19 to 50 years, the upper limit for iron intake is 45 mg/ day. Meanwhile zinc RDA for men is 11mg /day and 8 mg/day for females (Whiteney and Rolfes, 2008) [14]. The results were described forecasting the intake of 100-gram harvest plus wheat bread and assuming the suitability of each harvest plus genotype for 100-gram bread consumption frequency per day (one, two, three or four bread/s of 100-gram bread per day). Moreover, we considered other food sources of iron daily consumed and taken care in recommendation for not passing the upper limits of 45 mg/day for iron and 40 mg/ day for zinc.

7HPYT9, 7HPYT14, 8HPAN128, 8HPAN1, 8HPAN167, 8HPAN121,7HPYT49, 8HPAN8, 8HPAN19, 8HPAN31 and 7HPYT44 were having iron contents for two times consumption of 100-gram harvest plus bread. Meanwhile 8HPAN200 and 8HPAN42 iron contents were recommended for three times and four times consumption of the 100-gram bio fortified bread not exceeding the upper limit for iron intake.

Table 1: zinc, iron, phytate and tannin contents of harvest plus wheat samples

No.	Harvest Plus wheat	Zinc (mg/100g)	Iron(mg/100g)	Phytate (µg/g)	Tannin (mg/g)
1	7HPYT9	0.244a	14.1 a	0.189 a	0.151 a
2	7HPYT14	0.196 b	15.3 b	1.351 b	0.104 b
3	8HPAN150	0.322c	18.4 c	0.432 c	0.088 c
4	8HPAN200	0.57 d	10.54 d	0.459 d	0.086 d
5	8HPAN128	0.61 e	15.88 e	0.811 e	0.104 e
6	8HPAN177	0.856 f	9.84 f	0.459 f	0.098 f
7	8HPAN1	0.724 g	13.84 g	0.514 g	0.112 g
8	8HPAN17	0.758h	18.52 h	0.432 h	0.118 h
9	8HPAN42	0.908 i	8.6 i	0.946 i	0.114 i
10	8HPAN192	0.952 j	30.72 j	0.27 j	0.114 j
11	8HPAN159	1.048 k	31.3 k	0.378 k	0.116 k
12	8HPAN167	1.156l	12.38 l	0.595 l	0.122 l
13	7HPYT43	1.174 m	21.68 m	0.162 m	0.111 m
14	7HPYT12	1.182 n	17.98 n	0.108 n	0.132 n
15	8HPAN101	1.212 o	25.44 o	0.135 o	0.13 o
16	8HPAN121	1.218 p	16.94 p	0.378 p	0.124 p
17	7HPYT49	1.25 q	14.76 q	0.162 q	0.124 q
18	8HPAN62	1.358 r	21.6 r	0.351 r	0.128 r
19	8HPAN100	1.46 s	50.68 s	0.162 s	0.177 s
20	8HPAN8	1.552 t	14.78 t	1.108 t	0.139 t
21	8HPAN189	1.816 u	27.18 u	0.432 u	0.133 u
22	8HPAN19	1.634 v	16.26 v	0.027 v	0.118 v
23	7HPAN31	1.97 w	15.64 w	0.703 w	0.1 w
24	7HPYT41	1.916 x	18.94 x	0.784 x	0.108 x
25	8HPAN95	2.044 y	73.02 y	0.946 y	0.118 y
26	7HPYT44	2.29 z	17.36 z	0.108 z	0.1 z
27	7HPYT46	2.338 aa	23.54 aa	0.189 aa	0.106 aa
28	8HPAN173	2.402 ab	43.32 ab	0.649 ab	0.098 ab
29	7HPYT20	2.486 ac	26.4 ac	0.378 ac	0.094 ac
30	8HPAN50	2.736 ad	22.7 ad	0.865 ad	0.137 ad
31	7HPYT23	2.8616 ae	43.54 ae	0.676 ae	0.124 ae
32	8HPAN98	2.89 af	34.52 af	0.622 af	0.108 af
33	8HPAN76	3.95 ag	29.74 ag	0.405 ag	0.11 ag
34	7HPYT39	4.08ah	33.58 ah	0.649 ah	0.084 ah
35	8HPAN47	2.58 ai	38.78 aj	0.541 ai	0.086 ai
36	8HPAN191	7.134 aj	31.68 ai	0.27 aj	0.104 aj
37	8HPAN74	7.356 ak	89.1 ak	0.135 ak	0.096 ak
38	8HPAN11	12.232 al	37.4 al	0.324 al	0.086 al
39	7HPYT45	12.464 am	25.14 am	0.297 am	0.082 am
40	8HPAN29	22.414 an	27.26 an	0.243 an	0.112 an
41	8HPAN174	22.7 ao	39.48 ao	0.297 ao	0.139 ap
42	8HPAN43	26.632 ap	61.94 ap	0.162 ap	0.147 ao
43	7HPYT25	27.234 aq	30.02 aq	0.432 aq	0.141 aq
44	7HPYT4	46.398 ar	84.68 ar	0.595 ar	0.124 ar
45	8HPAN73	61.16 as	89.88 as	0.324 as	0.116 as
46	8HPAN94	66.44 at	39.84 at	0.757 at	0.132 at
	Minimum	0.20	8.60	0.03	0.08
	Maximum	62.74	89.88	1.35	0.18
	Grand mean	7.94	30.50	0.47	0.11
	Standard deviation	14.86	20.31	0.29	0.02

Means followed by different letters are significant at $p < 0.05$




Fig 1: Bar chart for iron and zinc content of harvest plus samples

	iron	zinc	phytate
zinc	0.622	0.000	
phytate	-0.086	0.004	
tannin	0.107	0.165	-0.091
	0.312	0.117	0.389
Cell Contents: Pearson correlation			
P-Value			

Fig 2: Correlation association among iron, zinc, phytate and tannin

As shown in the above figure 2 zinc and iron have high correlation coefficient in the harvest plus samples and it is in agreement with Zhao *et al.* (2009) [16] in which among the 150 lines of bread wheat, both grain Fe and Zn concentrations correlated positively and significantly (Ficco *et al.*, 2009) [2], their increment rate together in the sample provides good opportunity in genetic bio fortication of the harvest plus samples in bread wheat crop breeding as male or female parent and need only focus on the adoptability and stability of bio fortified bread wheat in different agro ecological zones and seasons.

Hailu Gebre-Mariam *et al.* (1991) [5] described the iron content of Ethiopian whole wheat grain being 1.5 mg/ 100 gm and Shahzad *et al.* (2014) stated that the iron and zinc concentration in whole grain of wheat are in the range of 2.9 to 7.3 mg/100 g and 0.7 to 8.5 mg/100 g, respectively and Wang *et al.* (2015) [13] stated zinc content of whole wheat grain to be 2 to 3.5 mg/ 100 g and that of wheat flour is about 1.5 mg/100g but the harvest plus samples analyzed in this study contained a 46 sample grand mean iron content of 30.50 mg/100 g which paves the way in cossombating Iron Deficiency Anaemia (IDA) and alleviate the impact of micronutrient deficiency on physical, mental and overall productivity of a society. The variation among wheat genotypes

in zinc concentration is described to be 1.5 to 4 times differences in genotypes (Wang *et al.* (2015) [13]. The difference in variation in iron and zinc content in germplasm and/or cultivar is known to be two to three-fold, iron content between 4.14 to 6.77 mg/ 100 g and zinc content ranging between 3.64 to 7.38 mg/ 100 g.

Kaur *et al.* (2020) [6] analysed 50 harvest plus genotypes from CIMMYT using energy dispersive x-ray fluorescence spectroscopy (EDXRF) instrument and found zinc content between 2.48 mg/100 g to 4.373 mg/100 g and Zhao *et al.* (2009) [16] found the zinc contents to be in the range of 1.35 mg/100 g to 3.45 mg /100 g, it is in close agreement with the 46 sample grand mean of 7.94 mg/100g but the iron content found by Kaur *et al.* from 50 samples was 2.96 mg/100 g to 3.766 mg/100g and Graham *et al.* (2007) reported the iron content of the bio fortified wheat was an average of 3.5 mg/100 g, and Wozniak and Makarski (2013) [15] stated iron content of biofortified wheat in their study was 2.468 mg/100 g to 2.62 mg/100 g; not in agreement with this study as the iron contents of the 46 samples was higher with a grand mean of 30.50 mg /100 g. Kaur *et al.* (2020) [6] noted that those harvest plus genotypes with both high iron and zinc content showed significant decrease in grain yield. Wang *et al.* (2015) [13] described the phytate content of whole wheat grain to be between 6 and 10 g/ kg and that of whole wheat flour is 1.5 to 3 g/ kg, but the 46 harvest plus whole wheat samples had phytate in the range of 0.03 to 0.47 µg/g meanwhile the phytic acid value found in this study was much lower than that reported by Wang *et al.* (2015) [13] and the lower phytate content is beneficial in lowering the phytic acid zinc molar ratio in higher zinc genotypes and consequently impacts zinc bio availability and it is noted that the phytic acid (PA)/Zn zinc molar ratio less than 15 is known to be important in improved zinc bioavailability, however, the phytic acid zinc molar ratio of not bio fortified whole wheat and wheat flour are found to be greater than 15 and thus low bioavailability of zinc (Wang *et al.*, 2015) [13]. Meanwhile Mackown *et al.* (2008) [7] described the condensed tannin of wheat in epicatechin equivalent condensed tannin with a mean of 0.343mg/g for experimental lines but the tannin concentration in the 46 sample was grand mean of 0.11 mg/g and it is in agreement with Mackown *et al.* (2008) [7] and the lower tannin concentration

is important in improving iron bioavailability from bio-fortified wheat.

Both Iron and zinc harvest plus genotypes had shown significant difference ($p < 0.05$) and 8HPAN100, 8HPAN95, 8HPAN74, 8HPAN43, 7HPYT4 and 8HPAN73 iron contents were above the upper level limit of 45 mg/day for adults, and consuming only one bread of 100 gram of those six samples stated above may cause gastro-intestinal distress, and the iron over load results in infections, fatigue, joint pain, skin pigmentation and organ damage (Whitney and Rolfes, 2008) [14].

Most of the harvest plus wheat samples had shown zinc content much lower than upper intake level per day except 7HPYT4, 8HPAN73, and 8HPAN94 above the upper limit of 40 mg/day and consuming one 100 gram of those breads of the three samples stated above may cause loss of appetite, impaired immunity, loss of High Density Lipoprotein (HDL), copper and iron deficiency (Whitney and Rolfes, 2008) [14].

But four harvest plus wheat samples 8HPAN121 (Fe, 16.94 mg/100g; Zn, 1.218 mg/100g), 8HPAN19 (Fe, 16.26 mg/100g; Zn, 1.634 mg/100g), 7HPAN31 (Fe, 15.64 mg/100g; Zn, 1.97 mg/100g) and 7HPYT44 (Fe, 17.36 mg/100g; Zn, 2.29 mg/100g) had shown promising results to be a better candidate for parental cross breeding since the iron contents shown to be safer to be consumed twice per day per 100 gram bread considering other sources of iron in day to day food consumed, relatively better zinc content and lower phytate contents.

Conclusion

Nutrition plays a key role in improving quality of agricultural produce. Hence crop breeding using genetic bio-fortification is best option as it reaches every household using the seed for cultivation specially bread wheat seed is most common cultivated in different parts of the country, thus iron and zinc bio-fortified wheat will reach every household within the baked bread, meanwhile the appropriate level of the micro-nutrients iron (Fe) and zinc (Zn) with respect to upper limit Recommended Daily Allowance (RDA) values per day without any possible effects of overdose intake along with the anti-nutritional factors (phytate and tannin) were assessed in the laboratory and 8HPAN121 (Fe, 16.94 mg/100g; Zn, 1.218 mg/100g), 8HPAN19 (Fe, 16.26 mg/100g; Zn, 1.634 mg/100g), 7HPAN31 (Fe, 15.64 mg/100g; Zn, 1.97 mg/100g) and 7HPYT44 (Fe, 17.36 mg/100g; Zn, 2.29 mg/100g) had shown promising results since the iron contents had shown to be safer to be consumed twice per day per 100 gram bread considering other sources of iron in day to day food consumed, relatively better zinc content and lower phytate contents consequently be a better candidate for parental cross breeding, adaptability and stability studies in different agro-ecological zones and seasons.

References

1. Bulbula DD, Uрга K. Study on the effect of traditional processing methods on nutritional composition and anti-nutritional factors in chickpea (*Cicer arietinum*). *Cogent Food & Agriculture*, 2018, 4.
2. Ficco DBM, Riefolo C, Nicastro G, De Simone V, Di Gesu AM, Beleggia R, *et al.* Phytate and mineral elements concentration in a collection of Italian durum wheat cultivars. *Field Crops Research*. 2009; 111:235-242.
3. Graham RD, Rosser JM. Carotenoids in staple foods: Their potential to improve human nutrition. *Food and Nutrition Bulletin*. 2000; 21:404-409.
4. Goudia BD, Hash CT. Breeding for high grain Fe and Zn levels in cereals. *International Journal of Innovation and Applied Studies*. 2015; 12(2):342-354.
5. Hailu Gebre-Mariam, Tanner DG, Mengistu Hulluka. eds. *Wheat Research in Ethiopia: A historical perspective*. Addis Ababa: IAR/CIMMYT, 1991.
6. Kaur N, Kaur H, Mavi GS. Assessment of nutritional and quality traits in biofortified bread wheat genotypes. *Food Chemistry*, 2020, 302.
7. MacKown CT, Carver BF, Edwards TJT. Occurrence of Condensed Tannins in Wheat and Feasibility for Reducing Pasture Bloat. *Crop Science*, 2008, 48.
8. Menaleshoa M, Sorsa G. Manual for soil, plant and water Analytical methods. JIJE analytical services, Addis Ababa, Ethiopia. Unpublished manual, 2010.
9. Monasterio I, Graham RD. Breeding for trace minerals in wheat. *Food and Nutrition Bulletin*, 2000; 21(4).
10. Oury FX, Leenhardt F, Remesy C, Chanliaud E, Duperrier B, Balfourier F, *et al.* Genetic variability and stability of grain magnesium, zinc and iron concentrations in bread wheat. *European Journal of Agronomy*. 2006; 25:177-185.
11. Shahzad Z, Rouached H, Rakha A. Combating Mineral Malnutrition through Iron and Zinc Biofortification of Cereals. *Comprehensive Reviews in Food Science and Food Safety*, 2014, 14.
12. Velu G, Singh RP, Huerta-Espino J, Peña RJ, Arun B, Mahendru-Singh A, *et al.* Performance of bio-fortified spring wheat genotypes in target environments for grain zinc and iron concentrations *Field Crops Research*. 2012; 137:261-267.
13. Wang Z, Liu Q, Pan F, Yuan L, Yin X. Effect of increasing rates of zinc fertilization on phytic acid /zinc molar ratio in zinc biofortified wheat. *Field Crops Research*. 2015; 184:58-64.
14. Whitney E, Rolfes SR. *Understanding nutrition*. 11th edition, Thomson Wadsworth, USA, 2018.
15. Wozniak A, Makarski B. Content of minerals, total protein and wet gluten in grain of spring wheat depending on cropping systems. *Journal of Elementology*, 2013.
16. Zhao FJ, Su YH, Dunhama SJ, Rakszegi M, Bedo Z, McGrath SP, *et al.* Variation in mineral micronutrient concentrations in grain of wheat lines of diverse origin. *Journal of Cereal Science*. 2009; 49:290-295.