



ISSN Print: 2664-6064
ISSN Online: 2664-6072
Impact Factor: RJIF 5.2
IJAN 2023; 5(1): 07-11
www.agriculturejournal.net
Received: 08-10-2022
Accepted: 12-11-2022

Arun Gangey
M.Sc Ag, Plant Physiology,
College of Agriculture,
Indira Gandhi Krishi
Vishwavidyalaya, Raipur,
Chhattisgarh, India

Dharmendra Khokhar
Scientist, Plant Physiology,
College of Agriculture,
Indira Gandhi Krishi
Vishwavidyalaya, Raipur,
Chhattisgarh, India

Corresponding Author:
Arun Gangey
M.Sc Ag, Plant Physiology,
College of Agriculture,
Indira Gandhi Krishi
Vishwavidyalaya, Raipur,
Chhattisgarh, India

Effect of growth promoter on morpho-physiological, biochemical parameters and essential oil yield of mentha (*Mentha arvensis* L.)

Arun Gangey and Dharmendra Khokhar

DOI: <https://doi.org/10.33545/26646064.2023.v5.i1a.90>

Abstract

A field experiment entitled “Effect of growth promoter on morpho-physiological, biochemical parameters and essential oil yield of mentha (*Mentha arvensis* L.)” The field experiment was carried out at Department of Plant Physiology, Agriculture Biochemistry and Medicinal and Aromatic Plants and Instructional Herbal Garden, IGKV, Raipur (C.G.) in *Mentha arvensis* L. during the *Rabi* season of 2020-21. Randomized block design (RBD) was used to conduct the experiment with three replications. There was 10 treatments T₁ (Control, Water Spray), T₂ (IAA @ 50 ppm), T₃ (IAA @ 70 ppm), T₄ (IAA @ 90 ppm), T₅ (GA3 @ 50 ppm), T₆ (GA3 @ 70 ppm), T₇ (GA3 @ 90 ppm), T₈ (BAP @ 50 ppm), T₉ (BAP @ 70 ppm), T₁₀ (BAP @ 90 ppm).

Experimental findings revealed that Herbage yield ranged from 18.07- 20.92 ton ha⁻¹. The highest herbage yield was recorded in T₈ (20.92 tonha⁻¹). Herbage yield in T₅ (20.88 tonha⁻¹) was statistically at par with T₈ and the minimum herbage yield was observed in T₁ (18.07 ton ha⁻¹). Essential oil (%) ranged from 0.64-0.76%. The highest essential oil (%) was obtained in T₆ (0.76%) which was at par with T₉ (0.73%), and the lowest essential oil (%) was observed in T₁ (0.64%).

Keywords: Mentha, PGR, essential oil

Introduction

Mentha spp. L. is known as mint belongs to the family Lamiaceae. It is important essential oil bearing plant. Mentha cultivation has gain economic importance due to its essential oil. The major component of the mentha essential oil is menthol, which is used in good, cosmetics and pharmaceuticals. India being largest producer of the world in mentha oil share 80-85% of world production. China, Brazil, US and Japan are the other major mentha oil producing countries. India being the leader in production of menthe oil, exports around 20,000-21,000 tons annually and maintain 1st position in the mentha oil world market. Farmers of Chhattisgarh have started the cultivation of mentha in few pockets of Bastar region. The crop may be an alternate crop to the summer paddy.

Materials and Methods

A field experiment entitled “Effect of growth promoter on morpho-physiological, biochemical parameters and essential oil yield of mentha (*Mentha arvensis* L.)” The field experiment was carried out at Department of Plant Physiology, Agriculture Biochemistry and Medicinal and Aromatic Plants and Instructional Herbal Garden, IGKV, Raipur (C.G.) in *Mentha arvensis* L. during the *Rabi* season of 2020-21. Raipur is located in the central area, agro-climatic zone of Chhattisgarh at latitude 21.6° N and longitude 81.36° E with an altitude of 289.56’ meters above the mean sea level. Raipur comes under dry, sub humid agro climatic region. The source of rainfall is south west monsoon. This region generally receives monsoon during June to September with annual rainfall of 1200-1400 mm and very little during October and February. The region had extreme temperature, May is the hottest and December is the coolest month of the year. The pattern of rainfall, particularly during June to September months has a great variation from year to year. The maximum temperature at Raipur goes up to 48 °C in summer while minimum temperature falls down up to 6 °C in winter season.

In this region the relative humidity is high from June to October, and wind velocity is high from May to August with its peak in June - July months.

The research trial was conducted in Randomized block design (RBD) There was 10 treatments T₁ (Control, Water Spray), T₂ (IAA @ 50 ppm), T₃ (IAA@70ppm), T₄ (IAA@ 90 ppm), T₅ (GA₃ @ 50 ppm), T₆(GA₃ @ 70 ppm), T₇(GA₃ @ 90 ppm), T₈ (BAP @ 50 ppm), T₉ (BAP @ 70 ppm), T₁₀ (BAP @ 90 ppm). and replicated twice. Nursery was raised in pro tray Coco peat and FYM were used as media for growing mentha cutting. The transplanting was done on 25th February, 2021. Transplanting were done as per requirement, healthy seedlings were transplanted with the flexible wire rope marked at specific interval (15x45 cm) with off colored cloth strips for maintaining spacing of (15x45 cm). Transplanting was done manually.

Five plants were selected as representative plant and tagged in each plot for recording growth as well as yield attributes. These plants were harvested separately for post harvest observations and herbage yield and oil content were later added corresponding total yield per plot. Harvesting was done with the help of sickles by manual labors

All data obtained in the was statistically analyzed using *F*-test, the procedure given by Gomez & Gomez (1984), critical difference (CD) values at *P*= 0.05 were used to determine the significance of differences between means.

Result and Discussion

Growth Parameters

The observations were recorded before the treatment and 15 days after the treatment. Recorded data was analyzed using the standard statistical tools. From the results it can be inferred that the plant height ranged from 37-58 cm. The maximum plant height was observed in T₇ (58.00 cm) which was statistically similar with T₆ (55.33 cm), and the minimum plant height was observed in T₁ (37 cm). Leaves plant⁻¹ ranged from 59.61-96.07. The number of leaves per plant maximum was observed in T₈ (96.07) which was statistically similar with T₅ (95.93), the number of leaves per plant minimum was observed in T₁ (59.61). Number of branches per plant was ranged from 4.45-7.80. The number of branches per plant was observed maximum in T₈ (7.80) which was statistically similar with T₅ (7.27), and the number of branches plant⁻¹ was observed minimum in T₁ (4.45). Fresh weight Plant⁻¹ ranged from 6.80-11.33 gm. The

highest fresh weight plant⁻¹ was recorded in T₈ (11.32 gm) which was at par with T₁₀ (10.71 gm), and the lowest fresh weight plant⁻¹ was recorded in T₁ (6.80 gm). Dry weight plant⁻¹ ranged from 2.16-3.73 gm. The highest dry weight plant⁻¹ was observed in T₈ (3.73 gm) which was at par with T₂ (3.16 gm), and the lowest dry weight plant⁻¹ was observed in T₁ (2.16 gm). Leaf area plant⁻¹ was ranged from 267.65-612.04 cm². The maximum leaf area plant⁻¹ was recorded in T₈ (612.04 cm²) which was at par with T₅ (552.12 cm²), and the minimum leaf area plant⁻¹ was recorded in T₁ (267.65 cm²) Leaf area index ranged from 2.78-5.43. The maximum leaf area index was recorded in T₈ (5.43) which was at par with T₂ (4.79) and T₄ (4.79). The minimum leaf area index was recorded in T₁ (2.78). Specific leaf area ranged from 264.73-548.44. The maximum specific leaf area was recorded in T₈ (548.44) which was at par with T₃ (539.10), and the minimum specific leaf area was recorded in T₁ (264.73). Specific leaf weight ranged from 0.19-0.35. The maximum specific leaf weight was recorded in T₈ (0.35) which was at par with T₄ (0.33), and the minimum specific leaf weight was recorded in T₁ (0.19). Total chlorophyll content ranged from 0.84-1.09 (mg g⁻¹ fresh wt.). The maximum total chlorophyll content was recorded in T₆ (1.09) which was at par with T₉ (1.05), and the minimum total chlorophyll was recorded in T₁ (0.84).

Herbage yield ranged from 18.07- 20.92 ton ha⁻¹. The highest herbage yield was recorded in T₈ (20.92 tonha⁻¹). Herbage yield in T₅ (20.88 tonha⁻¹) was statistically at par with T₈ and the minimum herbage yield was observed in T₁ (18.07ton ha⁻¹). Essential oil (%) ranged from 0.64-0.76%. The highest essential oil (%) was obtained in T₆ (0.76%) which was at par with T₉ (0.73%), and the lowest essential oil (%) was observed in T₁ (0.64%).

Yield attributes

Herbage yield ranged from 18.07- 20.92ton ha⁻¹. The highest herbage yield was recorded in T₈ (20.92 tonha⁻¹). Herbage yield in T₅ (20.88 tonha⁻¹) was statistically at par with T₈ and the minimum herbage yield was observed in T₁ (18.07ton ha⁻¹). Essential oil (%) ranged from 0.64-0.76%. The highest essential oil (%) was obtained in T₆ (0.76%) which was at par with T₉ (0.73%), and the lowest essential oil (%) was observed in T₁ (0.64%), Similar result was found by (Singh *et al.*, 1999^[48]).

Table 1: Effect of growth promoter on growth attributes of Mentha (*Mentha arvensis* L.).

Treatments	Growth attributes							
	Pre treatment				Post treatment			
	Plant height (cm)	Number of leaves plant ⁻¹	Number of branches plant ⁻¹	Fresh weight Plant ⁻¹ (g)	Plant height (cm)	Number of leaves plant ⁻¹	Number of branches plant ⁻¹	Fresh weight Plant ⁻¹ (g)
T ₁ Control	35.20	51.02	3.90	1.25	37.00	59.61	4.45	6.80
T ₂ IAA @ 50 ppm	35.37	47.03	3.93	2.90	39.00	69.93	6.87	9.81
T ₃ IAA @ 70 ppm	35.63	47.60	3.93	3.12	43.00	76.84	6.83	9.35
T ₄ IAA @ 90 ppm	33.40	48.70	3.90	3.24	38.67	86.33	6.97	10.03
T ₅ GA ₃ @ 50 ppm	34.63	48.92	3.87	3.78	54.33	95.93	7.27	10.42
T ₆ GA ₃ @ 70 ppm	34.37	47.83	3.90	3.89	55.33	63.43	7.10	9.33
T ₇ GA ₃ @ 90 ppm	34.13	47.09	3.87	3.01	58.00	66.93	6.47	8.55
T ₈ BAP @ 50 ppm	34.60	49.52	3.80	3.67	44.00	96.07	7.80	11.32
T ₉ BAP @ 70 ppm	34.87	47.00	3.77	3.90	41.00	70.83	4.73	10.29
T ₁₀ BAP @ 90 ppm	35.33	48.16	4.00	3.67	40.67	63.87	5.77	10.71
Sem±	1.78	1.51	0.10	0.94	3.59	7.72	0.71	0.84
CD=0.05%	NS	NS	NS	NS	10.67	22.93	2.12	2.51

Table 2: Effect of growth promoter on growth attributes of Mentha (*Mentha arvensis* L.).

Treatments	Growth attributes							
	Pre treatment				Post treatment			
	Dry weight plant ⁻¹ (g)	Leaf Area plant ⁻¹ (cm ²)	Leaf area index	SLA) (cm ² /gm)	Dry weight plant ⁻¹ (g)	Leaf Area plant ⁻¹ (cm ²)	Leaf area index	SLA) (cm ² /gm)
T ₁ Control	0.95	159.92	1.82	190.83	2.16	267.65	2.78	264.73
T ₂ IAA @ 50 ppm	1.12	157.97	1.90	291.69	3.16	500.32	4.79	446.48
T ₃ IAA @ 70 ppm	1.06	163.75	1.77	223.71	2.79	512.67	5.01	539.10
T ₄ IAA @ 90 ppm	1.00	165.71	1.70	258.84	2.56	529.40	4.79	478.12
T ₅ GA ₃ @ 50 ppm	1.13	154.61	1.75	301.43	2.47	552.12	3.19	398.41
T ₆ GA ₃ @ 70 ppm	1.11	162.30	1.79	195.42	2.39	437.82	3.00	360.37
T ₇ GA ₃ @ 90 ppm	1.01	162.10	1.73	186.38	2.26	485.67	3.54	446.46
T ₈ BAP @ 50 ppm	1.30	160.99	1.71	228.04	3.73	612.04	5.43	548.44
T ₉ BAP @ 70 ppm	1.08	160.94	1.85	247.22	2.36	464.51	2.85	441.01
T ₁₀ BAP @ 90 ppm	1.02	169.05	1.79	267.15	2.45	449.50	3.16	385.87
Sem±	0.03	1.3	0.03	5	0.37	6.0	0.53	50.9
CD=0.05%	NS	NS	NS	NS	1.10	8.4	0.56	151.5

Table 3: Effect of growth promoter on biochemical attributes of Mentha (*Mentha arvensis* L.).

Treatments	Biochemical attributes					
	Pre treatment			Post treatment		
	Chlorophyll content			Chlorophyll content		
	Chlorophyll a	Chlorophyll b	Total chlorophyll	Chlorophyll a	Chlorophyll b	Total chlorophyll
T ₁ Control	0.61	0.19	0.8	0.64	0.20	0.84
T ₂ IAA @ 50 ppm	0.59	0.22	0.81	0.79	0.22	1.01
T ₃ IAA @ 70 ppm	0.64	0.35	0.99	0.74	0.21	0.95
T ₄ IAA @ 90 ppm	0.62	0.34	0.96	0.68	0.21	0.89
T ₅ GA ₃ @ 50 ppm	0.61	0.26	0.87	0.71	0.21	0.92
T ₆ GA ₃ @ 70 ppm	0.65	0.26	0.91	0.83	0.29	1.12
T ₇ GA ₃ @ 90 ppm	0.63	0.34	0.97	0.80	0.22	1.02
T ₈ BAP @ 50 ppm	0.64	0.26	0.9	0.76	0.21	0.97
T ₉ BAP @ 70 ppm	0.55	0.24	0.79	0.82	0.23	1.05
T ₁₀ BAP @ 90 ppm	0.62	0.29	0.91	0.69	0.21	0.90
Sem±	0.04	0.06	0.27	0.03	0.01	0.04
CD=0.05%	NS	NS	NS	0.10	0.03	0.12

Table 4: Effect of growth promoter on growth attributes of Mentha (*Mentha arvensis* L.).

treatment	Herbage yield (kg/plot)	Herbage yield (t ha ⁻¹)	Essential oil (%)	Essential oil kg/ha
T ₁ Control	4.07	18.07	0.64	112.00
T ₂ IAA @ 50 ppm	4.53	20.14	0.70	125.52
T ₃ IAA @ 70 ppm	4.60	20.44	0.67	122.67
T ₄ IAA @ 90 ppm	4.67	20.73	0.65	112.15
T ₅ GA ₃ @ 50 ppm	4.70	20.88	0.66	120.04
T ₆ GA ₃ @ 70 ppm	4.20	18.66	0.76	146.26
T ₇ GA ₃ @ 90 ppm	4.37	19.40	0.71	132.14
T ₈ BAP @ 50 ppm	4.93	20.92	0.69	125.34
T ₉ BAP @ 70 ppm	4.53	20.16	0.73	139.87
T ₁₀ BAP @ 90 ppm	4.23	18.81	0.66	119.86
Sem±	0.08	0.36	0.08	3.20
CD=0.05%	0.25	1.07	0.25	9.50

Conclusions

The experiment was conducted first time at Instructional Herbal Garden hence, any concrete conclusion cannot be drawn from the data of one year experiment; however the following important inference may be drawn from the results of present investigation:

It is concluded from the study that T₈ (BAP @ 50 ppm) is the best treatment for morpho-physiological growth of mentha crop for getting higher herbage yield Treatment T₆ (GA₃ @ 70 ppm) is found to be superior for obtaining higher essential oil yield and economic value.

References

- Ahmad B, Jaleel H, Shabbir A, Khan MMA, Sadiq Y. Concomitant application of depolymerized chitosan and

GA₃ modulates photosynthesis, essential oil and menthol production in peppermint (*Mentha piperita* L.). Scientia Horticulturae. 2019;246:371-379.

- Akhtar N, Abdul Matinsarker M, Akhter H, Katrunnada M. Effect of planting time and micronutrient as zinc chloride on the growth, yield and oil content of *Mentha piperita*. Bangladesh journal of Scientific and Industrial Research. 2009;44(1):125-130.
- Alkire BH, Simon JE. Response of mid western peppermint (*Mentha X piperita* L.) and native Spearmint (*M. spicata* L.) to rate and form of nitrogen fertilizer. Acta Horticult. 1996;426:537-550.
- Arafa MM, El-Howeity MA. Effect of humic acid, plant growth promoting and methods of application on two potatoes (*Solanum tuberosum* L.) cultivar grown under

- sandy soil condition. *Menoufia Journal of Soil Science*. 2017;2(2):91-104.
5. Askary M, Talebi SM, Amini F, Bengal ADB. Effect of NaCl and iron oxide nanoparticles on *Mentha piperita* essential oil composition. *Environmental and Experimental Biology*. 2016;14:27-32.
 6. Ayyobi H, Peyvast GA, Olfati JA. Effect of vermicompost and vermicompost extract on oil yield and quality of peppermint (*Mentha piperita* L.). *Journal of Agricultural Sciences (Belgrade)*. 2013;58(1):51-60.
 7. Crysagryris A, Petropoulos AS, Fernades A, Barros NTF, Isabel CFR. Effect of phosphorus application rate on *Mentha spicata* L. growth in deep flow technique (DFT), *Food Chemistry*. 2019;276: 84-92.
 8. Deepak BR, Padma M, Prabhakarand BN, Vijaya D. Effect of NPK, zinc and sulphur levels on oil yield of Japanese mint (*Mentha arvensis* L.) var. Kosi. *Journal of Pharmacognosy and Phytochemistry*. 2019;8(4):3412-3415.
 9. Deepak BR, Padma M, Prabhakarand BN, Vijaya D. Effect of NPK, zinc and sulphur levels on oil yield of Japanese mint (*Mentha arvensis* L.) var. Kosi. *Journal of Pharmacognosy and Phytochemistry*. 2019;8(4):3412-3415.
 10. Duriyaprapan S, Britten EJ. The effect of age and location of leaf on quantity and quality of Japanese mint oil production. *Journal of experimental Botany*. 1982;33(4):810-814.
 11. Fathi A, Oveysi M, Nasri M, Tohidi H, Kasraei P. Physiological responses of peppermint (*Mentha piperita* L.) to plant growth regulators and salinity stress. *Iranian Journal of Plant Physiology*. 2020;11(1):3499-3508.
 12. Hamid AM, Reza AP, Saeid H, Morad S. Changes In Yield and Growth of Green Mint (*Mentha spicata* L.) Under Foliar Application of Urea and Soil Application of Vermicompost. *International journal of Advanced Biological and Biomedical Research*. 2014;2:2072-2076.
 13. Hendawy Saber. Effect of NP fertilization on Growth and Essential Oil of Lovage plants under Egyptian Conditions. *Middle East Journal of Agriculture Research*. 2014;3(4):1031-1036.
 14. Ivanova K, Manov M, Iliev L, Stefanov B. To Effect of Cytokinins on the Essential Oil Composition of *in Vitro* Produced Peppermint Plants. *Biotechnology & Biotechnological Equipment*; c1996.
 15. Izhar M, Khan M, Yasmin T, Zahid NY. Differential effect of fertilizers on menthol contents in mint (*Mentha arvensis* L.). *American Research Journal of Agriculture*. 2015;1(1):55-60.
 16. Juárez-Rosete CR, Olivo-Rivas A, Aguilar-Castillo JA, Bugarín-Montoya RUBEN, Arrieta-Ramos BG. Nutrition assessment of NPK in mint (*Mentha spicata* L.) cultivated in soilless system. *Annual Research & Review in Biology*; c2014. p. 2462-2470.
 17. Kapri M, Singh AK, Sisodia A, Padhi M. Influence of GA3 and BA (Benzyladenine) on flowering and post-harvest parameters in lily. *Journal of Pharmacognosy and Phytochemistry*. 2018;7(3):1916-1918.
 18. Kaymak HC, Yarali F, Guvenc I, Donmez MF. The effect of inoculation with plant growth rhizobacteria (PGPR) on root formation of mint (*Mentha piperita* L.) cuttings. *African journal of Biotechnology*, 2008, 7(24).
 19. Khanam D, Mohammad F. Effect of structurally different plant growth regulators (PGRs) on the concentration. Yield and constituents of peppermint essential oil. *Journal of Herb, science and medicinal plants*; c2016.
 20. Khanam Daraksha, Mohammad Firoz. Effect of Structurally Different Plant Growth Regulators (PGRs) on the Concentration, Yield, and Constituents of Peppermint Essential Oil. *Journal of Herbs, Spices & Medicinal Plants*. 2016;23:1-10.
 21. Khorasaninejad S, Mousavi A, Soltanloo H, Hemmati K, Khalighi A. The effect of salinity stress on growth parameters, essential oil yield and constituent of peppermint (*Mentha piperita* L.). *World Applied Sciences Journal*. 2010;11(11):1403-1407.
 22. Kumar P, Singh A, Lashiram N, Pandey RK, Dogra S, Iqbal J. Effect of plant growth regulator s on quality flower and seed production of marigold (*Tagetes erecta*) Bangladesh J. Bot. 2012;49(3):567-577,
 23. Kumar A, Mouzzam H, Singh P. Response Of Gibberellic Acid On Growth Behaviour and Menthol Oil Yield of Mentha (*Mentha piperita* L.) Hort Flora Research Spectrum. 2012;1(3):284-285.
 24. Kumar A, Patro HK Kewalanand. Effect of Zinc and Sulphur on herb, oil yield and quality of Menthol mint (*Mentha arvensis* L.) var. Kosi. *J Chem. Pharm. Res*. 2010;2(4):642-648.
 25. Lafmejani ZN, Jafari AA, Moradi P, Moghdam AL. Impact of foliar application of copper sulphate and copper nanoparticle on some morpho- physiological traits and essential oil composition of peppermint (*Mentha piperita*) *Herba Rolonica*. 2018;64(2):13-23.
 26. Lyczko J, Piotrowski K, Kolasa K, Galek R, Szumny A. *Mentha piperita* L. micropropagation and the potential influence of plant growth regulators on volatile organic compound composition. *Molecules*. 2020;25(11):2652.
 27. Mahantesh PS, Gangadharappa PM, Hiremath JS, Sharatbabu AG, Pooja MR, Nishchitha M. Effect of Row Spacing and Nitrogen Levels on Growth and Yield of Japanese Mint (*Mentha arvensis* L). *International Journal of Agriculture Sciences*. 2017;9(21):4304-4307.
 28. Mahmoodabad HA, Parchin RA, Hokmalipoor S, Shaban M. Changes in yield and growth of green Mint (*Mentha spicata* L.) Under foliar application of urea and soil application of vermicompost. *Int. J Adv. Biol. Biomed. Res*. 2014;2:2072-2076.
 29. Marino S, Ahmad U, Ferreira MI, Alvino A. Evaluation of the effect of irrigation on biometric growth, physiological response, and essential oil of (*Mentha spicata* L.). *Water*. 2019;11(11):2264.
 30. Mengel K, Kirkby EA. Principles of plant nutrition. 4th ed. *International Potash, Institute Switzerland*; c1987.
 31. Murray MT. The healing power of herbs: the enlightened person's guide to the wonders of medicinal plants. *Rocklin, CA: Prima Pub*. 1995;XIV:410.
 32. Nostro N, Germano MP, Angelo VD, Marino A, Cannatelli MA. Extraction methods and bioautography for evaluation of medicinal plant antimicrobial activity. *Lett Appl Microbiol*. 2000;30:379-384.
 33. Ormancey X, Sisalli S, Coutiere P. Formulation of essential oils in functional perfumery. *Parf Cosm Act*. 2001;157:30-40.
 34. Pande P, Chand S, Pandey A, Patra DD. Effect of sole and conjoint application of iron and manganese on herb

- yield, nutrient uptake, oil quality vis-à-vis their optional level in spearmint (*Mentha spicata* Linn. Emend. Nathh. C.v. Arka), Indian journal of natural products and Resources. 2011;2(2):242-249.
35. Prasad FM, Prakash V, Jillani AA, Prasad S, Thomas Arnold R, Robert KS. Effect of Naphthalene acetic acid (NAA) as growth regulator along with chelated zinc and iron on the availability of manganese, zinc, copper and iron in *Mentha piperita* Linn cultivar kukrail, International Journal of Environment. 2014;3(2):338.
 36. Rahman KM, Sattar MA, Rahman GMM. Effect of fertilizer and manures on growth and yield of Tulsi and Pudina Medicinal Plant. Journal of Environmental Science and Natural Resources. 2014;7(2):13-16.
 37. Roodbari N, Roodbari S, Ganjali A, Ansarifar M. The effect of salinity stress on growth parameters and essential oil percentage of peppermint (*Mentha piperita* L.). International Journal of Advanced Biological and Biomedical Research. 2013;1(9):1009-1015.
 38. Santoro VM, Nievas FL, Zygadlo JA, Giordano WF, Banchio E. Effects of growth regulators on biomass and the production of secondary metabolites in peppermint (*Mentha piperita* L.) micropropagated *in vitro*; c2013.
 39. Sawan ZM, Hafez SA, Basyony AE. Effect of phosphorus fertilization and foliar application of chelated zinc and calcium on seed, protein and oil yields and oil properties of cotton. J Agric Sci. 2001;136:191-198.
 40. Sawan Z, Hafez SA, Basyony AE. Effect of phosphorus fertilization and foliar application of chelated zinc and calcium on seed, protein and oil yields and oil properties of cotton. The Journal of Agricultural Science. 2001;136:191-198.
 41. Shekhar I, Daniel S, Kishore P, Singh AK, Kanaujia A. Impact of fertilizer dose and spacing on oil yield of Japanese mint (*Mentha arvensis* L.) var. Sarj. J Pharmacogn Phytochem. 2018;7(4):2463-2465.
 42. Souza MP, Matos NEO, Matos FJA. Constituintes Químicos de Plantas Medicinais Brasileiras. Imprensa Universitária/UFC, Fortaleza, 1991, 416.
 43. Suresh V, Fetricia JP, Saranya V, Sarithra S, Tamilselvan K. To study the effect of fym, coirpith, vermicompost, humic acid and panchagavya on growth and yield of mint (*Mentha arvensis* L.). Horticult Int J. 2018;2(6):417-419.
 44. Tucker AO. The truth about mints. Herb Companion. 1992;4:51-52.
 45. Valmorbida J, Boaro CS. Growth and development of *Mentha piperita* L. in nutrient solution as affected by rates of potassium. Brazilian Archives of Biology and Technology - Braz Arch Biol Technol, 2007, 50.
 46. Verma P, Pratapsingh P, Singh VR, Rahman L. Effect of combine doses of nitrogen and phosphorus on plant growth and oil yield of menthol mint (*Mentha arvensis* L.) in western Himalayan region of Utrakhand, International journal of herbal medicine. 2017;5(6):18-21.
 47. Verma PPS, Padalia RC, Singh VR. Effect of combined doses of nitrogen and phosphorus on plant growth and oil yield of menthol mint (*Mentha arvensis* L.) in western Himalayan region of Utrakhand.
 48. Singh B, Al-Haddad K, Chandra A. A review of active filters for power quality improvement. IEEE transactions on industrial electronics. 1999 Oct;46(5):960-971.